



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

so that the water may spout through it to the greatest horizontal distance on the plane. What difference in the result when the cylinder stands on a shelf of known height above the plane?

SOLUTION BY J. B. SMITH, Hampden-Sidney, Va.

Let a be the height of the cylinder and b the height of the orifice above the plane. Let v_0 be the velocity of the jet at the orifice. Then

$$v_0 = \sqrt{2g(a-b)}, \quad x_0 = v_0 t, \quad \text{and} \quad y = b - \frac{1}{2}gt^2.$$

Hence,

$$y = b - \frac{x^2}{4(a-b)}$$

is the equation of the trajectory, the origin being at the base of the cylinder and the y -axis being the vertical through the orifice.

For $y = 0$, we have $x^2 = 4b(a-b)$. As x^2 is a maximum when x is, we have

$$\frac{d(x^2)}{db} = 4(a-b) - 4b = 0,$$

$$\therefore b = a/2.$$

Hence the range will be a maximum when the height of the orifice is half that of the cylinder.

If the cylinder stands on a shelf of height h above the plane, we wish to find the maximum range on the line $y = -h$. In this case we have

$$x^2 = 4(b+h)(a-b).$$

It follows that the range is a maximum when $b = (a-h)/2$. If $h = a$, the orifice must be at the bottom of the cylinder; if $h > a$, b would be negative, but as this is impossible, the greatest range for the given cylinder is obtained by making the orifice at the bottom.

Also solved by HORACE OLSON and B. LIBBY.

NUMBER THEORY.

200. Proposed by R. D. CARMICHAEL, Indiana University.

Find the general solution, in relatively prime integers, of the equation $x^2 + y^2 = z^4$.

SOLUTION BY ARTEMAS MARTIN, Washington, D. C.

Let $x = p^2 - q^2$, $y = 2pq$; then

$$x^2 + y^2 = (p^2 - q^2)^2 + (2pq)^2 = (p^2 + q^2)^2 = z^4.$$

Let $p = r^2 - s^2$, $q = 2rs$; then

$$p^2 + q^2 = (r^2 - s^2)^2 + (2rs)^2 = (r^2 + s^2)^2;$$

therefore

$$x^2 + y^2 = (p^2 + q^2)^2 = (r^2 + s^2)^4 = z^4.$$

Retracing,

$$x = (r^2 - s^2)^2 - (2rs)^2, \quad y = 4rs(r^2 - s^2), \quad z = r^2 + s^2;$$

therefore

$$[(r^2 - s^2)^2 - (2rs)^2]^2 + [4rs(r^2 - s^2)]^2 = (r^2 + s^2)^4,$$

where r and s may be any numbers prime to each other, one odd and the other even.

Let $r = 2$, $s = 1$, and we have

$$7^2 + 24^2 = 625 = 25^2 = 5^4.$$

Let $r = 3$, $s = 2$; then we get

$$119^2 + 120^2 = 169^2 = 13^4.$$

Let $r = 4$, $s = 1$, and we find

$$161^2 + 240^2 = 289^2 = 17^4.$$

203. Proposed by R. D. CARMICHAEL, Indiana University.

Find solutions in integers of the equation

$$2x^2 + 1 = 3y^2. \quad (1)$$

I. SOLUTION BY E. E. WHITFORD, New York City.

Let $2x = z$. Then

$$z^2 - 6y^2 = -2. \quad (2)$$

By inspection the solution in smallest positive integers is $z_1 = 2$, $y_1 = 1$. The problem will not lose in generality if the solution be limited to positive integers. To represent concisely all the positive solutions without exception and without repetition I have derived the following formula:

$$z + y\sqrt{6} = \frac{(z_1 + y_1\sqrt{6})^{2k+1}}{2^k},$$

where

$$z_1 = 2, \quad y_1 = 1, \quad k = 0, 1, 2, 3, \dots$$

Therefore

$$z = 2, \quad 22, \quad 218, \quad 2158, \quad \dots,$$

$$y = 1, \quad 9, \quad 89, \quad 881, \quad \dots$$

Since the values of z are even each set gives a solution of equation (1).

$$x = 1, \quad 11, \quad 109, \quad 1079, \quad 10681, \quad \dots,$$

$$y = 1, \quad 9, \quad 89, \quad 881, \quad 8721, \quad \dots$$

These results were obtained by Euler in his "Algebra," by de la Roche in his "Larismetique" (1520), who copied from the "Triparty" of Chuquet (1484); and probably by nearly everyone who tried, before decimal fractions came into common use, to find the approximate value of the square root of 6.

Equation (2) is a slightly generalized form of the Pell equation $x^2 - Ay^2 = 1$.